



RESEARCH ARTICLE

**DETERMINATION OF ACID INSOLUBLE ASH VALUES OF SOME MEDICINAL PLANTS OF GENUS
*SESBANIA***Kadam V.B.¹, R.K.Momin², M.S.Wadikar³, S. B. Andhale⁴¹P.G.Department of Botany & Research Centre, K.T.H.M. College, Nasik – 422002²Department of Botany, Milliya Arts, Science and Management Science College, Beed³Department of Botany, Vinayakrao Patil Science College, Vaijapur, Aurangabad⁴Department of Botany, M.V.P.Samaj's Science College Dindori, Nashik

Received 20 August 2013; Revised 30 August 2013; Accepted 4 September 2013

ABSTRACT

The seasonal variation of total ash, acid soluble ash and acid insoluble ash have been investigated leaves, wood and bark of *Sesbania rostrata*, *Sesbania exaltata* and *Sesbania sesban*, which are medicinally important. Comparative account of total ash, acid soluble ash and acid insoluble ash content of bark of *Sesbania rostrata* showed high level of total ash (range 7.98 % to 8.70 %) and low level of total ash of leaves of *Sesbania sesban* (range 6.34 % to 6.78 %). The acid soluble ash showed higher level of bark of *Sesbania rostrata* (range 7.45 % to 7.74 %) and lower in wood of *Sesbania sesban* (range 2.62 % to 2.83 %). Comparative account of acid insoluble ash of bark of *Sesbania rostrata* showed higher (range 1.91 % to 2.24 %) and lower in the wood of *Sesbania sesban* (range 0.37 % to 0.58 %).

Key Words: Total ash, acid soluble ash, acid insoluble ash, medicinal plant, *Sesbania*

INTRODUCTION:

The phytochemical constituents and medicinal properties of most of the medicinal plants were recorded in the last few decades by a number of workers (Nudrat and Usha, 2005). Medicinal plants have been used as traditional treatments for numerous human diseases for thousands of years. Medicinal properties of plants are due to the active chemical constituents present in different parts of the plant (Mitscher et al, 1980). Medicinal plants continue to be an important therapeutic aid for the ailments of humankind. The survey and documentation of medicinally important plants in each and every place is very much important for easy identification of local traditional healers, conservation and sustainable utilization. All human beings require a number of complex organic/inorganic compounds in diet to meet the need for their activities. The important constituents of diet are carbohydrates, fats, proteins, vitamins, minerals and water (Indrayan et al., 2005). Plants are the rich source of all the elements essential for human beings. The quality of many foods depends on the concentration and type of minerals what they contains, also play a very significant role against a variety of degenerative diseases and processes, they may also prevent and reduce injury from environmental pollutant and enhance the ability to work and learn. It is clear that mineral nutrition is important to maintain good health.

The use of mineral element is found to have been developed and used widely to cure several health problems. The amount and composition of ash remaining after combustion of plant material varies considerably according to the part of the plant, age, treatment etc. The constituents of the ash also vary with time and from organ to organ. Ash usually represents the inorganic part of the plant (Tambe and Kadam, 2012).

Today, there is a renewed interest in traditional medicine and increasing demands for more drugs from plant sources. This revival of interest in plant-derived drugs is mainly due to the current widespread belief that "greenmedicine" is safe and more dependable than the costly synthetic drugs, many of which have adverse side effects. Nature has bestowed upon us a very rich botanical wealth and a large number of diverse types of plants grow wild in different parts of our country. The seasonal variation of water soluble extractive alcohol soluble extractive and ether soluble extractive have been investigated in leaves, wood and bark of *Sesbania rostrata*, *Sesbania exaltata* and *Sesbania sesban*.

Sesbania rostrata as an important dietary nutritive source in Southeast Asian country's. *Sesbania rostrata* are richest source of amino acid, minerals and antioxidants vitamins. This species is unique because it fixes nitrogen not only in its roots in the soil, but also in its aerial parts including stems and branches (Dutt et.al., 1983). Various

parts of this plant are used in Indian traditional medicine for the treatment of diuretic, emetic, fevers, headaches, anemia, bronchitis, inflammation, leprosy, gout, rheumatism, anxiolytic, anticonvulsive and hepatoprotective (Pari and Uma, 2003). It also has anti-inflammatory, analgesic and antipyretic activity (Momin and Kadam, 2011).

Sesbania exaltata is a crop generally cultivated for its nutritive value to soil. It is cultivated in monsoon season almost throughout India and grows sandy, loamy and clay soils. It is an ideal green manure crop as it is quick-growing, succulent, and easily decomposable with low moisture requirements and produces maximum amount of organic matter and nitrogen in the soil. Seed flour is used in the treatment of ringworm, skin diseases and wounds. The mature seeds of this species are known to be cooked and eaten by the Indian tribal's (Brown, 1954).

Sesbania sesban seeds considered stimulants and astringent. Leaves considered purgative, anthelmintic and anti-inflammatory. Leaves showed a high crude protein content, 25 to 30% and are a useful source of protein for ruminant diets and a source of supplement fodder for livestock. Study of the effect of *Sesbania sesban* seed powder on female albino rats showed inhibition of ovarian function, change of uterine structure and prevention of implantation with 100 % control of fertility (Shiv Pal Singh, 1990). The aqueous extracts of leaves in STZ-induced diabetic rats showed significant increase in serum insulin and HDL level and decreases in blood glucose, total cholesterol and triglycerides when compared to glibenclamide (Pandhare et.al.,2011). *Sesbania sesban* was referred to as milk shrub. Farmers were encouraged to feed *Sesbania* fodder to lactating cows to enhance milk secretion (Brown, 1954).

MATERIAL AND METHODS:

Method recommended in pharmacopoeia of India (Anonymous, 1966), and British Pharmacopoeia (Anonymous, 1973) were followed for determining ash value and percentage method.

***Preparation of Ash-** 3gm of drug was incinerated in a Silica crucible over the burner. The charred material was heated in muffle furnace for six hours at 600-650 °C. The ash formed was white and free from carbon. It was cooled and weighed on the ash less filter paper.

***Determination of acid insoluble ash-** The acid was boiled for 5 minutes with 25ml of dilute hydrochloric acid. Insoluble matter collected in crucible or on an ash less filter Paper and washed with hot water, ignited and

weight. Percentage of acid insoluble ash was calculated with reference to the air dried drug.

RESULT AND DISCUSSION:

Sesbania rostrata – The total ash of leaves range from (7.98 % to 8.70 %) higher level of total ash show at summer (8.70 %) than monsoon (7.98 %) and winter (8.55 %). Total ash of bark showing higher level at summer 9.98 % as compared to monsoon 9.36 % and winter 9.69 %. Total ash of wood content show higher level at summer 4.45 % than winter 4.07 % and monsoon 3.88 %. The percentage of total ash were found to be in the increasing order of bark < leaves < wood. The range of acid soluble ash content of leaves was ranging from (6.35 % to 6.73 %) among different season tested (Table). Wood show lower level of acid soluble ash content (range from 2.97 % to 3.30 %), higher level show at summer 3.30 % than winter 3.10 % and monsoon 2.97 % . Bark show higher level of acid soluble ash at summer 7.74 % as compared to monsoon 7.45 % and winter 7.56 %. The percentage of water soluble ash were found to be in the increasing order of bark < leaves < wood. Acid insoluble ash of leaves was higher at summer 1.97 % as compared to monsoon 1.63 % and winter 1.85 %. The range of percentage of acid insoluble ash of bark show higher (1.91 % to 2.24 %) than leaves (1.63 % to 1.97 %) and wood (0.91 % to 1.15 %). In wood acid insoluble ash show higher level at summer 1.15 % than winter 0.97 % and monsoon 0.91 %, while in bark highest acid insolubility of ash at summer 2.24 % than winter 2.13 % and monsoon 1.91 %. The percentage of acid insoluble ash content were found to be in increasing order of bark < leaves < wood (Table.1).

Sesbania exaltata - The percentage of total ash content of leaves show highest level at summer 7.67 % as compared to monsoon 6.78 % and winter 7.43 %. In bark total ash content observed higher level at summer 8.97 % as compared to winter 8.56 % and monsoon 8.32 % , while in wood total ash rages from (3.45 % to 3.96 %) highest level observed at summer 3.96 % than monsoon 3.45 % and winter 3.67 %. The percentage of total ash content were found to be in increasing order of bark < leaves < wood. The acid soluble ash of leaves was measures in summer (5.82 %), monsoon (5.13 %) and winter (5.65 %). The acid soluble ash of bark was highest at summer (7.12 %) than winter (6.91 %) and monsoon (6.78 %). The acid soluble ash of wood ranges from (2.83 % to 3.07 %) , higher being in summer 3.07 % as compared to monsoon 2.83 % and winter 3.01 % . The percentage of acid soluble ash were found to be in the increasing order of bark < leaves < wood. The acid insoluble ash content of leaves was measure in summer

(1.85 %), monsoon (1.65 %) and winter (1.78 %). The bark show acid insoluble ash range from (1.54 % to 1.85 %). The higher level being observed in summer 1.85 % as compared to monsoon 1.54 % and winter 1.65 % , while in wood show low acid insolubility of ash, it measure at monsoon 0.62 % ,winter 0.66 % and summer 0.89 % . The percentage of water insolubility were found to be in the increasing order of bark < leaves < wood (Table.1).

Sesbania sesban - The total ash of leaves range from (6.34 % to 6.78 %) higher level of total ash show at summer (6.78 %) than monsoon (6.34 %) and winter (6.55 %). Total ash of bark showing higher level at summer 8.79 % as compared to monsoon 8.13 % and winter 8.37 % . Total ash of wood content show higher level at summer 3.41 % than winter 3.22 % and monsoon 2.99 % . The percentage of total ash were found to be in the increasing order of bark < leaves < wood. The range of acid soluble ash content of leaves was ranging from (4.83 % to 5.04 %) among different season tested (Table).

Wood show lower level of acid soluble ash content (range from 2.62 % to 2.83 %) , higher level show at summer 2.83 % than winter 2.66 % and monsoon 2.62 % . Bark show higher level of acid soluble ash at summer 6.93 % as compared to monsoon 6.78 % and winter 6.84 % . The percentage of acid solubility ash were found to be in the increasing order of bark < leaves < wood. Acid insoluble ash of leaves was higher at summer 1.74 % as compared to monsoon 1.51 % and winter 1.69 % . The range of percentage of acid insoluble ash of bark shows higher (41.35 % to 1.86 %) than leaves (1.51 % to 1.74 %) and wood (0.37 % to 0.58 %). In wood acid insoluble ash show higher level at summer 0.58 % than winter 0.56 % and monsoon 0.37 % , while in bark highest acid insolubility of ash at summer 1.86 % than winter 1.53 % and monsoon. 1.35 % . The percentage of acid insoluble ash content were found to be in increasing order of bark < leaves < wood (Table.1)

Table 1: Determination of Ash Values of *Sesbania rostrata*, *Sesbania exaltata* and *Sesbania sesban*

Plant parts	Season	Total ash (%)			Acid soluble (%)			Acid insoluble (%)		
		Plant 1	Plant 2	Plant 3	Plant 1	Plant 2	Plant 3	Plant 1	Plant 2	Plant 3
Leaves	Summer	8.70	7.67	6.78	6.73	5.82	5.04	1.97	1.85	1.74
	Monsoon	7.98	6.78	6.34	6.35	5.13	4.83	1.63	1.65	1.51
	Winter	8.55	7.43	6.55	6.70	5.65	4.86	1.85	1.78	1.69
Wood	Summer	4.45	3.96	3.41	3.30	3.07	2.83	1.15	0.89	0.58
	Monsoon	3.88	3.45	2.99	2.97	2.83	2.62	0.91	0.62	0.37
	Winter	4.07	3.67	3.22	3.10	3.01	2.66	0.97	0.66	0.56
Bark	Summer	9.98	8.97	8.79	7.74	7.12	6.93	2.24	1.85	1.86
	Monsoon	9.36	8.32	8.13	7.45	6.78	6.78	1.91	1.54	1.35
	Winter	9.69	8.56	8.37	7.56	6.91	6.84	2.13	1.65	1.53

Plant 1- *Sesbania rostrata*, Plant 2- *Sesbania exaltata* and Plant 3- *Sesbania sesban*

REFERENCES:

1. Anonymous, 1966: Pharmacopoeia of india. New Delhi, Government of India.
2. Anonymous, 1973: British Pharmacopoeia. By Her Majesty's stationary office, London, U.K
3. Brown, W.H. (1954) Useful plants of the Philippines. *Republic of Philippines Dept. of Agriculture and Natural Resources Technical Bulletin* No.10 ,Vol. 2
4. Dutt, A.K., Pathania, U. and Kumar, V. (1983) Growth of *Sesbania sesban*. Nitrogen Fixing Tree. *Research Report* 1, 5 - 6.
5. Indrayan, A.K., Sharma, S., Durgapal, D., Kumar, N. and Kumar, M., 2005: Determination of nutritive value and analysis of mineral elements for some medicinally valued plants from Uttaranchal. *Current Sci* 89, 1252-1255.
6. Kadam, V.B. (2009) "Determination of ash values of three endangered medicinal plants of South Gujarat Forest". *Plant Archives*, 9 (1), 27 – 29.
7. Mitscher, L.A., Park, Y.H. and Clark, D.1980: Antimicrobial agents from higher Research Article, 2, (2).
8. R.K.Momin and Kadam V.B.(2011) "Determination of ash values of Some Medicinal Plants of Genus *Sesbania* of Marathwada Region in Maharashtra" *Journal of Phytology* , 3(2),52-54.

9. Nudrat Z, Sayed and Usha Mukundan.2005. *Medicinal and aromatic plants of India* Part I, In: Khan and Khanum, Ukaaz Publications, Hyderabad
10. Pandhare, R.B., Sangameswaran, B., Mohite, P.B. and Khanage, S.G.(2011) Antidiabetic Activity of aqueous extract of *Sesbania sesban* in Streptozotocin induced Diabetic rats. *AJMB Archives*, 3, 1
11. Pari,L. and Uma,A. (2003) Protective effect of *Sesbania grandiflora* against Erythromycin estolate induced hepatotoxicity. *Therapie*, 58: 439-443
12. Shiv Pal Singh (1990) Fertility control of female through *Sesbania sesban* seeds. *The journal of research and education in Indian medicine*, 9 (4): 27 – 32.
13. Tambe S.S. and Kadam V.B (2010) “Determination of ash values of two endangered medicinal taxa of Marathwada Region.” *Journal of Ecobiotechnology* 2/8, 25-28.