



## Research Article

### EVALUATION OF ANTI-HYPERTENSIVE ACTIVITY OF NORDOSTACHYS JATAMANSION AGAINST CADMIUM CHLORIDE INDUCES HYPERTENSION IN ALBINO RATS

Bharat Kumar Tiwari<sup>1</sup>, Charanjeet Singh<sup>2</sup>, Anil Godara<sup>3</sup>

<sup>1</sup>M.Pharm. Research Scholar, Department of Pharmacology, Jaipur College of Pharmacy, Jaipur, Rajasthan, India

<sup>2</sup>Associate Professor, Department of Pharmacology, Jaipur College of Pharmacy, Jaipur, Rajasthan, India

<sup>3</sup>Lecturer, Department of Pharmacology, Jaipur College of Pharmacy, Jaipur, Rajasthan, India

Received 02 Dec. 2017; Accepted 20 Jan. 2018

#### ABSTRACT

*Nardostachys jatamansi* is a very effective, potential and safe drug for the management of patients with essential hypertension along with dietary restrictions and modified lifestyle. The sesquiterpenevaleranone isolated from the subterranean parts of *Nardostachys jatamansi* (DC) is found to having sedative, tranquilizing and antihypertensive properties. *Nardostachys jatamansi* whole plant was collected from herbal garden of National Institute of Ayurveda (NIA), Jaipur. Genuine sample of plant material was weighed by the electronic monopan balance and a thin layer of sample was spreaded on a white colour sheet. By bull lens, the layer was examined for foreign matter. Moisture content was determined by placing weighed sample of 5gm of Plant material in oven at 105°C for 5 hours, and calculated weight of sample for every 30 minute. The pH value of an aqueous liquid may be defined as the common reciprocal of the hydrogen ion concentration expressed in gram per liter Freshly prepared extracts were tested for the presence of various active phytochemicals like phenols, tannin, Flavonoids, protein, reducing sugar, carbohydrates, lipids, Saponin, triterpenoid alkaloid, resins, volatile oils, anthraquinone and Quinone. Identification can be effected by observation of spots of identical RF value. Acute Toxicity Study (according to OECD 423 guideline). B. Anti-hypertensive activity (CdCl<sub>2</sub> Induced hypertension)

**Keywords:** *Nardostachys jatamansi*, Acute Toxicity, hypertensive activity, Ayurveda

#### 1. INTRODUCTION:

Hypertension is a serious disease affecting a significant population globally The reported prevalence of hypertension varied around the world, with the lowest prevalence in rural India (3.4% in men and 6.8% in women) and the highest prevalence in Poland (68.9% in men and 72.5% in women). Awareness of hypertension was reported for 46% of the studies and varied from 25.2% in Korea to 75% in Barbados; treatment varied from

10.7% in Mexico to 66% in Barbados and control (blood pressure < 140/90 mmHg while on antihypertensive medication) varied from 5.4% in Korea to 58% in Barbados.

#### 1.1 Plant collection

*Nardostachys jatamansi* whole plant were collected from herbal garden of National Institute of Ayurveda (NIA), Jaipur and authenticated from CSIR-national institute of science communication and information resources, New Delhi.

Table 1: Table showing chemicals used during the dissertation work

Hydrochloric acid	Sodium hydroxide	Toluene
Distilled Water	Acetic acid	Hypophosphorus
Ethanol	Potassium chromate	Lead acetate
Petroleum ether	Potassium iodide	Potassium bismuth iodide
Potassium Ferro cyanide	Ammonia	Vanillin

Sulphuric acid	Ammonium chloride	Ferric chloride
Potassium thiocyanate	Sodium sulphide	Chloroform
Per chloric acid	Sodium potassium tartrate	Pyridine
Barium chloride	Iodine solution	Magnesium turning
Methanol	Copper sulphate	Petroleum ether
Molisch's reagent	Picric acid	Glycerine

## 1.2. Equipment's and Apparatus

List of equipment's and apparatus used during the dissertation work.

**Table 2: Table showing list of equipment's & apparatus.**

Digital balance	Condenser
Magnifying Bull Lens	Heating mantle
Hot air oven	Hot plate
Silica crucible	Water bath
Grinder	Common glass wares
Muffle furnace	Rotary shaker
Desiccators	

## 1.3. Physiochemical Properties

**a. Determination of foreign matter:** Genuine sample of plant material was weighed by the electronic monopan balance and a thin layer of sample was spreaded on a white colour sheet. By bull lens, the layer was examined for foreign matter

**b. Determination of moisture content:** Moisture content was determined by placing weighed sample of 5gm of Plant material in oven at 105°C for 5 hours, and calculated weight of sample for every 30 minute, until the weight of the sample came out to be constant, no variation of weight would be recorded.

**c. Determination of p:** HPractically it means the quantitative indication of the acidity or basic nature of a solution. The pH of a given solution is measured by using digital pH meter.

**d. Determination of Total Ash:** Silica Crucible was cleaned, dried well & labeled with glass pencils and then weighed to constant weight. 5 gm of powdered drug sample was put in the Silica crucible. The drug was spread evenly into a thin layer.

**e. Determination of Water Soluble Ash:** Water soluble ash value was determined as per Pharmacopoeia of India 1996. Boil the total ash for

5 minutes with 25 ml of water; collect the insoluble matter in a Gooch's.

**f. Determination of Extractive values:** Extraction of plant material is done as per API suggested method

**g. Determination of Alcohol Soluble Extractive:** It was taken Macerate 5 gm of the air dried drug, coarsely powdered, with 100 ml of alcohol the specified strength in a closed flask for twenty-four hours, shaking frequently during six hours and allow to stand for eighteen hours. Filter rapidly, taking precautions against loss of solvent, evaporate 25 ml of the filtrate to dryness in a tarred flat bottomed shallow dish, and dry at 105°C, to constant weight and weigh. Calculate the percentage of alcohol-soluble extractive with reference to the air-dried drug.

**h. Determination of Water Soluble Extractive:** Proceed as directed for the determination of alcohol-soluble extractive, using distilled water instead of ethanol.

**i. Determination of Petroleum Ether Soluble Extractive (Fixed Oil Content):** Transferred a suitably weighed quantity (depending on the fixed oil content) of the air-dried, crushed drug to an extraction thimble, extract with solvent ether (or petroleum ether, b.p. 40°C to 60°C) in a continuous extraction apparatus for 6 hours. Filtered the

extract quantitatively into a tarred evaporating dish and evaporate off the solvent on a water bath. Dry the residue at 105° C to constant weight. Calculate the percentage of ether-soluble extractive with reference to the air-dried drug.

**1.4. Phytochemical Properties:** Freshly prepared extracts were tested for the presence of various active phytochemicals like phenols, tannin, Flavonoids, protein, reducing sugar, carbohydrates, lipids, Saponin, triterpenoid alkaloid, resins, volatile oils, anthraquinone and Quinone.

**a. Chromatography plates-**

T.L.C. plate coated with 0.25 mm layer of silica gel GF 254 with fluorescent indicator, (Mercks) were used. (Each plate dimension is 10 cm long and 2 cm width)

**b. Activation of pre-coated Silica gel G60F254** Dry in hot oven at 105° C for one to two hour.

**c. Preparation of Test sample:** Alcoholic Extract

**d. Preparation of mobile solution:** Toluene: Ethyl acetate 9.3:0.7. (v/v)

**e. Sample application:** Sample was applied with the help of capillary 1(one) cm above the base of T.L.C. plate. Then it was dipped in mobile solution. T.L.C. plate was removed from the mobile solution immediately after the spot reached the 1(one) cm below the top of the T.L.C. plate.

**f. Visualization:** AnisaldehydeSulphuric Acid

**g. Rf. Value:** Measure and record the distance of each spot from the point of its application and calculate the Rf. value by dividing the distance travelled by the spots by the distance travelled by the front of the mobile phase.


**Table 3: Physicochemical Properties**

Sr. No	Tests	Observation
1	Foreign matter	4.27%
2	Moisture content	9.34%
3	pH	5.6
4	Total Ash	7.35%
5	Water Soluble Ash	5.97%
6	Acid Insoluble Ash	2.47%
7	Alcohol Soluble Extractive	9.24%
8	Water Soluble Extractive	17.31%
9	Petroleum Ether Soluble Extractive	6.34%

**Table 4: Phytochemical Properties**

S. No	Name of Test	Observation
1	Tests for Carbohydrates-	
A	Molisch's Test	+ve
B	Benedict's test	-ve
C	Barfoed's test	-ve
D	Fehling solution test	+ve
2	Tests for Alkaloids-	
A	Mayer's reagent test	+ve
B	Dragondroff's reagent test	-ve
C	Wagner's test	+ve
D	Hager's Test	-ve
3	Test for Amino acids-	
A	Ninhydrin test	+ve
4	Protein	
A	Xanthoprotic test	+ve

B	Millons test	-ve
C	Biuret test	+ve
5	Tests for Glycosides-	
a	Borntrager's Test	
6	Test for Phenolic Compound:	+ve
7	Test for Saponin(Foam test)-	+ve
8	Test for Steroids( Salkowaskireaction)	+ve
9	Test for Tennins –	
A	FeCl <sub>3</sub>	-ve
B	Lead acetate	+ve
c	Pot. Dichromate	-ve

**TLC:**Fig. 1: TLCPlate of *Nardostachysjatamansi*R<sub>f</sub> Value= 0.73, 0.82, 0.86, 0.97**1.5 PHARMACOLOGICAL STUDY****1.5.1 Pre-Clinical Study:**

A. Acute Toxicity Study (according to OECD 423 guideline).

B. Anti-hypertensive activity (CdCl<sub>2</sub> Induced hypertension)

- **Ethical clearance:** The Experimental study was conducted on Albino Wistarrats after approval from the Institutional Animal Ethical Committee. Experimental study was conducted in Institute of Biomedical and Industrial Research, Vidyadhar Nagar, Jaipur. We had taken IAEC approval from ethical committee of institute of biomedical and industrial research. CPCSEA Approval No: 1737/PO /RC/14/CPCSEA, IAEC No:ibir/cpcsea/iaec/2017 /7/15.

**Methods:**

- **Preparation of aqueous extract: (Test drug)**

Dried plant material was ground to coarse powder 20gm. drug was extracted with 200ml. of water using Soxhlet apparatus (Hot extraction method) till 72 hrs. At the end of extraction the extract was concentrated and solvent was totally evaporated in a petridish with the help of water bath. The thick and sticky paste thus obtained was stored in air tight container at room temperature till further use. And this extract was used as test drug.

**A. Oral Acute Toxicity Study**

Acute toxicity was done according to OECD guideline 423 ANNEX 2c

**Anti-hypertensive activity:- CdCl<sub>2</sub>Induced hypertension in albino wistar rats**

A total of 60 albino wistarrats (both sex) were used in 4 month to 6 month age range and having weight about 70-140gm.in this study. These were divided into 5 groups of 6 rats each in both model. They were kept in separate cages after marking for 24hours prior to study. They were given food and water ad libitum.

**Marking of albino wistar rat for identification:**

The albino rat was marked with Picric acid in each group as H, B, T, HB, BT and HT where:-

- H stands for Head of albino rat
- B stands for Back of albino rat
- T stands for Tail of albino rat

- HB stands for Head back of albino rat
- BT stands for Back Tail of albino rat
- HT stands for Head tail of albino rat.

- **Test drug dose Calculated for Experimental Albino Rats –**

Test dose was derived from Acute Toxicity Study.

**Table 5: Drug used for the Treatment of induced hypertension in Animal Groups**

Group	No. of rats	Intervention	Duration
A	6	1% CMC solution	4 week
B	6	Test drug (Aqueous Extract of whole plant of <i>Nardostachysjatamansi</i> )	4 week
C	6	Verapamil (15 mg/kg/day, P.O.)	4 week

The dose for experimental study was calculated by extrapolating the human dose to animal's dose based on the body surface area ratio. Hence the suitable dose for albino rat was calculated by referring to the table of **Paget and Barne's** i.e.

- ✓ Hypothetical Human dose of extract of formulation = 1000mg twice daily
- ✓ Conversion factor (Human to Albino rat) = 0.018
- ✓ 18 albino wistar rats of both sexes of 3 groups (A, B and C) were administered 1mg/kg/day Intra peritoneal (0.1% CdCl<sub>2</sub> solution) CdCl<sub>2</sub> for 15 days for inducing hypertension. On day 16<sup>th</sup> rats were weighed & their blood pressure was measured by non-invasive BP system for rodents and test sample had been administer for 30 days orally, after every week systolic blood pressure and diastolic blood pressure of each rats were measured.

Measurement using tail cuff method, rats were trained for at least one week until the BP was steadily recorded with minimal stress and restrain. Systolic and diastolic BP was measured weekly for 4 weeks by indirect non-invasive tail-cuff method.

#### 1.6 STATISTICAL ANALYSIS:

The observed data had been recorded and presented in tabular format. These data had been analyzed to establish patterns of results. The data had been subjected to ANOVA test to assess the statistical significance of the findings

**Acute oral toxicity study: (As per the OECD Guidelines 423 ANNEX 2c)-Acute Oral Toxicity (OECD GUIDELINE 423):-** Observation Report of test drug

**Table 6: Observations of toxicity at dose 2000mg/kg/body wt. of test drug**

Observation	30min.	4hr.	24hr.	48hr.	1week	2week
Skin and Fur	Normal	Normal	Normal	Normal	Normal	Normal
Eyes	Normal	Normal	Normal	Normal	Normal	Normal
Mucous Membrane	Normal	Normal	Normal	Normal	Normal	Normal
Salivation	Normal	Normal	Normal	Normal	Normal	Normal
Lethargy	Nil	Nil	Nil	Nil	Nil	Nil
Sleep	Normal	Normal	Normal	Normal	Normal	Normal
Coma	Nil	Nil	Nil	Nil	Nil	Nil
Convulsions	Nil	Nil	Nil	Nil	Nil	Nil
Tremors	Nil	Nil	Nil	Nil	Nil	Nil
Diarrhoea	Nil	Nil	Nil	Nil	Nil	Nil
Morbidity	Normal	Normal	Normal	Normal	Normal	Normal
Mortality	Nil	Nil	Nil	Nil	Nil	Nil

#### Hematological test:

Table 7: Hematological test report:

Sr. No.	Hematological Parameters	300 mg/kg	2000 mg/kg	Normal Range
1.	Haemoglobin	14.5	13.7	11.5-16.1
2.	WBC	6.9	6.4	6.6-12.6 x 10 <sup>3</sup> /mm <sup>3</sup>
3.	RBC	8.4	6.98	6.76-9.75 x 10 <sup>6</sup> / mm <sup>3</sup>
4.	Neutrophils	2.56	2.27	1.77-3.38 x10 <sup>3</sup> / mm <sup>3</sup>
5.	Lymphocytes	9.1	8.4	4.78-9.12 x 10 <sup>3</sup> / mm <sup>3</sup>
6.	Eosinophils	0.04	0.04	0.03-0.08 x 10 <sup>3</sup> / mm <sup>3</sup>
7.	Monocytes	0.02	0.01	0.01-0.04 x 10 <sup>3</sup> / mm <sup>3</sup>
8.	Basophiles	0.0	0.0	0.00-0.03 x 10 <sup>3</sup> / mm <sup>3</sup>
9.	Platelets	396	421	150-460 x 10 <sup>3</sup> /mol

**CdCl<sub>2</sub> induce Hypertension in Albino Rats**

Table 8: Diastolic Pressures of different groups at different time interval

Table 9: Systolic Pressure of different groups at different time interval

Systolic	1 day	8 day	15 day	23 day	30 day	37 day
	Mean± SEM	Mean± SEM	Mean± SEM	Mean± SEM	Mean± SEM	Mean± SEM
Group A	138±1.862	139.83±1.364	140.67±1.687	139.67±2.459	140.67±1.563	138.83±1.772
Group B	139.83±2.056	133±0.601	124.67±1.116	123.17±1.576	120.5±1.118	142±1.36
Group C	140.83±1.195	121.17±1.222	119.5±1.204	120.17±0.98	119.83±0.792	150.5±6.854

Diastolic	1 day	8 day	15 day	23 day	30 day	37 day
	Mean± SEM	Mean± SEM	Mean± SEM	Mean± SEM	Mean± SEM	Mean± SEM
Group A	108.67±2.246	109±2.671	111.17±2.496	111±3.13	106.5±1.057	107.83±1.922
Group B	107.17±1.352	93±1.125	84.33±1.282	82±0.816	81±0.856	107.33±1.745
Group C	103.5±2.432	81±0.966	79.83±0.601	78.5±0.992	81.17±1.138	106±1.713

Table 10: Comparison of hypertension between group A, group B & group C

Dunnett's multiple comparisons test	Mean Diff.	95.00% CI of diff.	Significant?	Summary	Adjusted P Value
8 day					
Group A vs. Group B	6.83	0.2238 to 13.44	Yes	*	0.0404
Group A vs. Group C	18.66	12.05 to 25.27	Yes	****	0.0001
15 day					
Group A vs. Group B	16	9.394 to 22.61	Yes	****	0.0001
Group A vs. Group C	21.17	14.56 to 27.78	Yes	****	0.0001
23 day					
Group A vs. Group B	16.5	9.894 to 23.11	Yes	****	0.0001
Group A vs. Group C	19.5	12.89 to 26.11	Yes	****	0.0001
30 day					
Group A vs. Group B	20.17	13.56 to 26.78	Yes	****	0.0001
Group A vs. Group C	20.84	14.23 to 27.45	Yes	****	0.0001
37 day					
Group A vs. Group B	-3.17	-9.776 to 3.436	No	ns	0.5801
ension Group A vs. Group C	-11.67	-18.28 to -5.064	Yes	****	0.0001

**REFERNCES:**

1. Arora RB, Arora CK, Sha MJ, Shet UK. Animal species variation in hypotensive activity Ofjatamansone with a report in the clinical trial of this drug. Indian J Med Sci 1967.
2. Hypertens,2004 Jan;22(1):11-9 [Worldwise prevalence of hypertension:a systemic review. Kearney PM:,wheltonM, ReynoldsK, ,Whelton PK,He J.
3. Ali M. Pharmacognosy and Phytochemistry, Vol. 1st; CBS Publisher and Distributors, New Delhi, 2008, 672-673.
4. Kokate CK, Prohit AP, Gokhale SB. Pharmacognosy, Edn46, Vol 1st ,2nd, Nirmal Prakshan, Pune, 2010, 1.48.
5. Evans WC. Trease and Evans Pharmacognosy, Edn 15,published by Elseveir; Noida, India, 2008
6. Chatterjee B, Basak U, Dutta J, Banerji A, NeumanT.Prange Studies on the Chemical Constituents of *N.jatamansi*Cheminform 2005; 36:17.
7. Bose BC, Vijayvarngiya R, BhatnagarJN. *Nardostachysjatamansi*DC: a phytochemical study of its active constituents. Indian J Med Science 1957; 11(10):799-802.
8. Ali M. Pharmacognosy and Phytochemistry, Vol. 1st; CBSPublisher and Distributors, New Delhi, 2008, 672-673.
9. Dravyaguna Vigyanam (Materia medica vegetable drugs) (English-Sanskrit) Part-I(A-J), by Prof.Dr.GyanendraPandey, Edition-3rd, Chaukhambha Krishnadas Academy Varanasi, 2005. P. 836-838.